

# DETERMINANTS OF ADJACENCY MATRICES OF GRAPHS

ALIREZA ABDOLLAHI

ABSTRACT. We study the set of all determinants of adjacency matrices of graphs with a given number of vertices.

## 1. Introduction

Let  $G$  be a simple graph with finite number of vertices. We denote by  $\det(G)$  the determinant of an adjacency matrix of  $G$ . This number  $\det(G)$  is an integer and is an invariant of  $G$  so that its value is independent of the choice of vertices in an adjacency matrix. In this paper, we study the distributions of  $\det(G)$  whenever  $G$  runs over graphs with finite  $n$  vertices for a given integer  $n \geq 1$ . We denote by  $\mathcal{G}_n$  the set of all non-isomorphic graphs with  $n$  vertices and

$$\mathcal{DG}_n = \{ \det(G) \mid G \in \mathcal{G}_n \}, \quad \alpha_n = \max \mathcal{DG}_n \quad \text{and} \quad \beta_n = \min \mathcal{DG}_n.$$

In [2], Fallat and van den Driessche studied, among others things, the maximum  $W(n, k)$  and minimum  $w(n, k)$  of non-zero absolute values of determinants of  $k$ -regular graphs with  $n$  vertices and they determined  $W(n, 2)$ ,  $W(n, n-3)$  and  $w(n, 2)$ ,  $w(n, n-3)$ .

It is a famous result due to Hadamard [4] that if  $A = [a_{ij}]$  is an  $n \times n$  complex matrix such that  $\|a_{ij}\| \leq \mu$  for all  $i, j$  then  $\|\det(A)\| \leq \mu^n n^{\frac{n}{2}}$ . In [8], Ryser found an upper bound for the absolute value of a  $(0, 1)$  square matrix of size  $n$  with  $t$  non-zero entries. One may use Ryser's result to give an upper bound for the absolute values of the determinants of graphs with  $n$  vertices and  $m$  edges (see Theorem 2.5, below).

Using Brendan McKay's nauty, we have computed  $\mathcal{DG}_n$  for all  $n \leq 9$  (see Proposition 2.2, below).

Newman [7, Theorem 2.2] proved that  $k \cdot \gcd(k, n)$  divides the determinants of  $n \times n$   $(0, 1)$  matrices whose all row and column sums equal to  $k$ . A similar result for the determinants of graphs is proved in Theorem 2.6.

Hu [5] has determined the determinant of graphs with exactly one cycle. Here we obtain the possible determinants of graphs with exactly two cycles (see Proposition 2.11, below).

## 2. Preliminaries

**Proposition 2.1.** *Let  $K_n$  and  $P_n$  be the complete graph and the path with  $n$  vertices, respectively, and  $C_m$  be the cycle with  $m \geq 3$  vertices.*

---

1991 *Mathematics Subject Classification.* 05C50; 15A15.

*Key words and phrases.* Determinant; adjacency matrices of graphs; maximum determinant;

$$\begin{aligned}
(1) \quad \det(K_n) &= (-1)^{n-1}(n-1). \\
(2) \quad \det(P_n) &= \begin{cases} 0 & \text{if } n \text{ is odd} \\ 1 & \text{if } n = 2k, k \text{ is even} \\ -1 & \text{if } n = 2k, k \text{ is odd} \end{cases} \\
(3) \quad \det(C_m) &= \begin{cases} 2 & \text{if } m \text{ is odd} \\ 0 & \text{if } m = 2k, k \text{ is even} \\ -4 & \text{if } m = 2k, k \text{ is odd} \end{cases}
\end{aligned}$$

*Proof.* (1) It is well-known.

(2) It follows from the fact that  $\det(P_n) = -\det(P_{n-2})$ .

(3) It follows from [1, Theorem 1.3, p. 32].  $\square$

**Proposition 2.2.** *The following table gives the set of all determinants of graphs with at most 9 vertices. By  $i^j$  in the  $n$ -th row of the table, we mean that there are exactly  $j$  non-isomorphic graphs with  $n$  vertices whose determinants are all equal to  $i$ , and a single number  $i$  (with no exponent) shows that there is a unique graph with  $n$  vertices whose determinant is equal to  $i$ .*

| Number of Vertices | Determinants of graphs with multiplicities                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|--------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1                  | 0                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| 2                  | 0, -1                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| 3                  | $0^3, 2$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| 4                  | $-3, 0^7, 1^3$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| 5                  | $-4, 0^{25}, 2^6, 4$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| 6                  | $-5^3, -4^5, -1^{32}, 0^{99}, 3^{10}, 4^2, 7^2$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| 7                  | $-12^2, -10^2, -6^{13}, -4^{21}, -2^{20}, 0^{690}, 2^{204}, 4^{40}, 6^{17}, 8^{25}, 10^5, 12^5$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| 8                  | $-28^2, -27^2, -24^5, -23^5, -20^7, -19^{21}, -16^{51}, -15^{43}, -12^{90}, -11^{79}, -8^{128},$<br>$-7^{251}, -4^{581}, -3^{813}, 0^{6551}, 1^{2416}, 4^{758}, 5^{240}, 8^{73}, 9^{139}, 12^{24}, 13^{23}, 16^{32}, 17^8, 20^1, 21^3$                                                                                                                                                                                                                                                                                                                                                                         |
| 9                  | $-128^2, -96^3, -72^{12}, -64^7, -60^5, -56^{17}, -54^{12}, -50^{27}, -48^{13}, -46^{20}, -44^{39}, -42^{47},$<br>$-40^{103}, -38^{52}, -36^{110}, -34^{128}, -32^{593}, -30^{199}, -28^{295}, -26^{392}, -24^{765}, -22^{579},$<br>$-20^{869}, -18^{2747}, -16^{2247}, -14^{1805}, -12^{3062}, -10^{4290}, -8^{17582}, -6^{8531}, -4^{14901},$<br>$-2^{57065}, 0^{133174}, 2^{6767}, 4^{6950}, 6^{4669}, 8^{1566}, 10^{1349}, 12^{1156}, 14^{695}, 16^{606},$<br>$18^{106}, 20^{297}, 22^{173}, 24^{240}, 26^{95}, 28^{91}, 30^{61}, 32^{46}, 34^5, 36^{32}, 38^{28}, 40^3, 42^{17}, 44^{16}, 54^3, 60^3, 64$ |

*Proof.* Using nauty of Brendan McKay, one can generate all graphs with at most 9 vertices and then by GAP for example, it is easy to find the determinants.  $\square$

Here are some questions which are motivated by the data given in Proposition 2.2.

**Proposition 2.3.** (1)  $\beta_n \neq 0$  if and only if  $\beta_n < 0$  if and only if  $n \notin \{1, 3\}$ .  
(2)  $\alpha_n \neq 0$  if and only if  $\alpha_n > 0$  if and only if  $n \notin \{1, 2\}$ .  
(3) If  $G$  is a graph with odd number of vertices, then  $\det(G)$  is even.  
(4) The determinant of every bipartite graph with odd vertices is zero.

*Proof.* (1) It is clear that if  $n = 1$  or  $3$ , then  $\beta_n = 0$  and if  $n = 2$ , then  $\beta_n = -1$ . Thus assume that  $n \geq 4$ . It is enough to show that  $\beta_n < 0$ . If  $n$  is odd, we have  $\beta_n \leq \det(C_{n-2} + P_2) = -2$ , by Lemma 2.1. If  $n$  is even, we have  $\beta_n \leq \det(K_n) = -(n-1)$ , by Lemma 2.1. This proves (2).

(2) If  $n = 1$  or  $2$ , then clearly  $\alpha_n = 0$ . Thus it is enough to prove that  $\alpha_n > 0$

whenever  $n \geq 3$ . If  $n$  is odd, then  $\alpha_n \geq \det(K_n) = n - 1$ , by Lemma 2.1. If  $n$  is even, then  $\alpha_n \geq \det(K_{n-2} + P_2) = n - 3$ . This completes the proof of (3).

(3) Let  $n = |V(G)|$  be odd and let  $\mathcal{F}$  be an elementary figure (if exists) of  $G$  with  $n$  vertices. Then, by definition,  $\mathcal{F}$  is a disjoint union of  $s$  number of edges and  $t$  number of cycles. If  $t = 0$ , then  $n = 2s$  which is not possible. It follows that any elementary figure of  $G$  with  $n$  vertices have at least one cycle. Now it follows from [1, Theorem 1.3, p. 32] that  $\det(G)$  is an even number.

(4) Let  $G$  be a bipartite graph with odd  $n$  vertices. Since every bipartite graph has no odd cycles, it follows that  $G$  has no elementary figure with  $n$  vertices. Again [1, Theorem 1.3, p. 32] implies that  $\det(G) = 0$ .  $\square$

**Question 2.4.** For a given positive integer  $n$ , find integers  $m$  such that  $\det(G) \neq m$  for all graphs  $G$  with  $n$  vertices.

**Theorem 2.5.** Let  $G$  be a graph with  $n \geq 2$  vertices and  $m$  edges. Then  $|\det(G)| \leq \left(\frac{2m}{n}\right)^n \left(1 - \frac{2m-n}{n(n-1)}\right)^{n-1}$ .

*Proof.* Note that the number of 1's in any adjacency matrix of  $G$  is equal to  $2m$ . Now the proof follows from Theorem 3 of [8].  $\square$

**Proposition 2.6.** Let  $G$  be a graph with  $n$  vertices and let  $\{d_1, \dots, d_n\}$  be the set of vertex degrees of  $G$ . If  $d = \gcd(d_1, \dots, d_n)$ , then  $d \cdot \gcd(\frac{2m}{d}, d)$  divides  $\det(G)$ .

*Proof.* Let  $A = [A_1 \ \cdots \ A_n]$  be an adjacency matrix of  $G$ , where  $A_1, \dots, A_n$  are its columns. Then

$$\det(G) = |A_1 + \cdots + A_n \ A_2 \ \cdots \ A_n| = |\mathbf{d} \ A_2 \ \cdots \ A_n|,$$

where  $\mathbf{d} = [d_1 \ \cdots \ d_n]^T$ . Now sum up all the rows with the  $n$ th one. Then  $\det(G)$  is equal to the determinant of a matrix whose first column is

$$[d_1 \ \cdots \ d_{n-1} \ \sum_{i=1}^n d_i]^T$$

and its  $n$ th row is

$$[\sum_{i=1}^n d_i \ d_2 \ \cdots \ d_n].$$

Since  $\sum_{i=1}^n d_i = 2m$ , by factoring  $d$  from the first column and  $\gcd(\frac{2m}{d}, d_2, \dots, d_n)$  from the last column, we have that  $\det(G) = d \cdot \gcd(\frac{2m}{d}, d_2, \dots, d_n)$ . Since  $d_1 = 2m - \sum_{i=2}^n d_i$ , we have  $\gcd(\frac{2m}{d}, d_2, \dots, d_n) = \gcd(\frac{2m}{d}, d)$ . This completes the proof.  $\square$

**Theorem 2.7.** (Hu [5]) Let  $G$  be a connected graph with  $n$  vertices having a unique cycle  $C$  with  $k < n$  vertices. Then

$$\det(G) = \begin{cases} 1 & G \text{ has a perfect matchig, } k \equiv 1 \pmod{2}, n \equiv 0 \pmod{4} \\ -1 & G \text{ has a perfect matchig, } k \equiv 1 \pmod{2}, n \equiv 2 \pmod{4} \\ 4 & G \text{ has a perfect matchig, } k \equiv 2 \pmod{4}, n \equiv 0 \pmod{4} \\ -4 & G \text{ has a perfect matchig, } k \equiv 2 \pmod{4}, n \equiv 2 \pmod{4} \\ 0 & G \text{ has a perfect matchig, } k \equiv 0 \pmod{4} \\ 2 & G \text{ has no perfect matchig, } G \setminus C \text{ has a perfect matching, } n - k \equiv 0 \pmod{4} \\ -2 & G \text{ has no perfect matchig, } G \setminus C \text{ has a perfect matching, } n - k \equiv 2 \pmod{4} \\ 0 & \text{both } G \text{ and } G \setminus C \text{ have no perfect matching} \end{cases}.$$

**Corollary 2.8.** *If  $\alpha_n = \det(G)$  or  $\beta_n = \det(H)$  for some graphs  $G$  and  $H$ , then neither  $G$  nor  $H$  have a unique cycle for all  $n > 5$ .*

*Proof.* It is not hard to see that for  $n > 5$ ,  $\alpha_n > 4$  and  $\beta_n < -4$ . Since a graph with a unique cycle has exactly one connected component with a unique cycle and the other components (if exist) are tree (whose determinants belong to  $\{0, 1, -1\}$ ), Theorem 2.7 completes the proof.  $\square$

**Lemma 2.9.** *Let  $G$  be a graph containing exactly two cycles  $C_1$  and  $C_2$  of orders  $k$  and  $\ell$ , respectively such that  $|V(C_1) \cap V(C_2)| = 1$ . If  $|V(G)| = k + \ell - 1$ , then*

$$\det(G) = \begin{cases} 0 & \text{if both } k, \ell \text{ are even} \\ 2 \cdot (-1)^{\ell+k-1} \cdot \left( (-1)^{\frac{\ell-1}{2}+1} + (-1)^{\frac{k-1}{2}+1} \right) & \text{if both } k, \ell \text{ are odd} \\ 2 \cdot (-1)^{\ell+k-1} \cdot \left( (-1)^{\frac{\ell}{2}+\frac{k-1}{2}} + (-1)^{\frac{k-1}{2}+1} \right) & \text{if } k \text{ is odd, } \ell \text{ is even} \\ 2 \cdot (-1)^{\ell+k-1} \cdot \left( (-1)^{\frac{k}{2}+\frac{\ell-1}{2}} + (-1)^{\frac{\ell-1}{2}+1} \right) & \text{if } \ell \text{ is odd, } k \text{ is even} \end{cases}$$

*Proof.* Let  $C_1 := x_1 \cdots x_k x_1$  and  $C_2 := y_1 \cdots y_\ell y_1$ . Suppose  $C_1 \cap C_2 = \{x_k\} = \{y_\ell\}$ . We want to apply [1, Theorem 1.3, p. 32] and so we need to find all elementary figures of  $G$ , that are all spanning subgraphs of  $G$  with exactly  $k + \ell - 1$  vertices whose connected components are either an edge or a cycle of  $G$ . If  $k$  and  $\ell$  are both even, then  $G$  has no elementary figure. If  $k$  and  $\ell$  are both odd, then we have only two elementary figures

$$F_1 = \{C_1, y_1 y_2, \dots, y_{\ell-2} y_\ell\} \text{ and } F_2 = \{C_2, x_1 x_2, \dots, x_{k-2} x_k\}.$$

If  $k$  is odd and  $\ell$  is even, we have exactly 3 elementary figures as follows:

$$\{y_1 y_2, \dots, y_{\ell-1} y_\ell, x_1 x_2, \dots, x_{k-1} x_k\}, \{y_1 y_\ell, y_2 y_3, \dots, y_{\ell-2} y_{\ell-1}, x_{k-1} x_k, \dots, x_2 x_1\}, \\ \{C_2, x_1 x_2, \dots, x_{k-2} x_k\}.$$

Similarly, for the case  $k$  even and  $\ell$  odd, we have exactly 3 elementary figures (interchange  $k$  and  $\ell$  in the latter elementary figures). Now we can apply [1, Theorem 1.3, p. 32] and this completes the proof.  $\square$

**Lemma 2.10.** *Let  $G$  be a connected graph containing exactly two cycles  $C_1$  and  $C_2$  of orders  $k$  and  $\ell$ , respectively such that  $V(C_1) \cap V(C_2) = \emptyset$ . If  $V(G) = V(C_1) \cup V(C_2)$ , then  $\det(G) \in \{-8, 0, 3, 5, 16\}$ . If  $C_1$  and  $C_2$  are connected by a path  $P$  of length at least 2 and  $V(G) = V(C_1) \cup V(C_2) \cup V(P)$ , then*

$$\det(G) \in \begin{cases} \{0, \pm 4\} & \text{if } t = 3 \\ \{-16, -8, -5, -3, 0\} & \text{if } t = 4 \\ \{0, \pm 3, \pm 4, \pm 5, \pm 8, \pm 16\} & \text{if } t > 4 \end{cases}.$$

*Proof.* Let  $C_1 = x_1 \cdots x_k x_1$ ,  $C_2 = y_1 \cdots y_\ell y_1$ . By hypothesis, in any case, we have that there exists a path  $P = x_k = z_1 z_2 \cdots z_t = y_\ell$  connecting  $C_1$  and  $C_2$  such that  $V(G) = V(C_1) \cup V(C_2) \cup V(P)$ . It is clear that  $t = 2$  if and only if  $V(G) = V(C_1) \cup V(C_2)$ . In this case, by [1, Theorem 2.12, p. 59], we have  $\det(G) = \det(C_1) \det(C_2) - \det(P_{k-1}) \det(P_{\ell-1})$ . Now it follows from Lemma 2.1,  $\det(G) \in \{-8, 0, 3, 5, 16\}$ .

Now assume that  $t \geq 3$ , that is the length of  $P$  is at least 3, and let  $H$  be the induced subgraph of  $G$  on the vertices  $V = V(C_1) \cup (V(P) \setminus \{z_t\})$ . Note that, by

[1, Theorem 2.11, p. 59], we have

$$\det(H) = \begin{cases} -\det(P_{k-1}) & \text{if } t = 3 \\ \det(C_k) \det(P_{t-2}) - \det(P_{k-1}) \det(P_{t-3}) & \text{if } t \geq 4 \end{cases}.$$

Thus it follows from [1, Theorem 2.12, p. 59] that

$$\det(G) = \det(H) \det(C_\ell) - \det(H \setminus \{z_{t-1}\}) \det(P_{\ell-1}).$$

Note that  $\det(H \setminus \{z_{t-1}\})$  can be similarly compute by a formulae as given for  $\det(H)$ . Now it is easy to complete the proof.  $\square$

**Proposition 2.11.** *Let  $G$  be a graph with exactly two cycles. Then  $\det(G) \in \{0, \pm 1, \pm 2, \pm 3, \pm 4, \pm 5, \pm 8, \pm 16\}$ .*

*Proof.* By using [1, Theorem 2.11, p. 59] on the (possible) vertices of  $G$  with degree 1, we find that there exist a (possibly empty) forest  $F$  and either a connected graph  $H$  as of the form in Lemmas 2.9 or 2.10 or two connected graphs  $H_1$  and  $H_2$  each of which contains exactly one cycles such that  $\det(G) = \pm \det(F) \cdot \det(H)$  or  $\det(G) = \pm \det(F) \cdot \det(H_1) \cdot \det(H_2)$ , respectively, where  $\det(F) = 1$  if  $F$  is empty. Now Lemmas 2.9 and 2.10 and Theorem 2.7 complete the proof.  $\square$

Let us end the paper by the following problems and questions mainly arising from the data table given in Proposition 2.2 and some other investigations.

### Problems.

- 1) Describe  $\mathcal{DG}_n = \{\det(G) \mid G \in \mathcal{G}_n\}$ .
- 2) What are the maximum  $\alpha_n$  and minimum  $\beta_n$ ?
- 3) For a given  $n$ , what integers can never belong to  $\mathcal{D}_n$ ?
- 4) Is it possible to determine graphs  $G, H \in \mathcal{G}_n$  such that  $\det(G) = \alpha_n$  and  $\det(H) = \beta_n$ ? Do they have some distinguished properties from other graphs with  $n$  vertices? For example, must they always be connected?
- 5) Find relations between  $\mathcal{DG}_n$  and  $\mathcal{DG}_{n+1}$ .
- 6) Is it true that  $\alpha_n < \alpha_{n+1}$  for all  $n > 3$ ?
- 7) Is it true that  $\beta_{n+1} < \beta_n$ ?
- 8) Is it true that  $|\beta_n| > \alpha_n$  for all  $n > 7$ ?
- 9) Does  $\delta_0 := \lim_{n \rightarrow \infty} \frac{|\{G \in \mathcal{G}_n \mid \det(G)=0\}|}{|\mathcal{G}_n|}$  exist? If so, is it true that  $\delta_0 = \frac{1}{2}$ ?

### REFERENCES

- [1] D. M. Cvetković, M. Doob, H. Sachs, Spectra of graphs. Theory and application. Pure and Applied Mathematics, 87. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1980.
- [2] S. Fallat and P. van den Driessche, Maximum determinant of  $(0, 1)$  matrices with certain constant row and column sums. Linear and Multilinear Algebra 42 (1997), no. 4, 303–318.
- [3] F. C. Lin, Singularity of the adjacency matrix of graphs with non-interactive bicycle. J. Math. Study 37 (2004), no. 3, 321–324.
- [4] J. Hadamard, Résolution d’une question relative aux determinants, Selecta, pp. 136–142, Gauthier-Villars, Paris, 1935 and Oeuvres, Tome I, pp. 239–245, CNRS, Paris, 1968.
- [5] S. Hu, The Classification and maximum determinants of the adjacency matrices of graphs with one cycle. J. Math. Study 36 (2003), no. 1, 102–104.
- [6] S. Hu, H. F. Yan, The maximum determinants of the adjacency matrix of graphs with  $k$  cycles. Math. Appl. (Wuhan) 16 (2003), suppl., 125–128.

- [7] M. Newman, Combinatorial matrices with small determinants, *Canad. J. Math.* 30 (1978), 756-762.
- [8] H.J. Ryser, Maximal determinants in combinatorial investigations, *Canad. J. Math.* 8 (1956), 245-249.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ISFAHAN, ISFAHAN 81746-73441, IRAN  
*E-mail address:* `a.abdollahi@math.ui.ac.ir`